

Mechanism-based pharmacodynamic modeling for predicting exposure response

Meindert Danhof (1,2), Oscar E. Della Pasqua (1) and Bart A. Ploeger(1,2)

(1) Leiden University, Leiden-Amsterdam Center for Drug Research, Division of Pharmacology, Leiden, the Netherlands. (2) LAP&P Consultants BV, Leiden, the Netherlands

Objectives: A major challenge in drug discovery and development is the prediction, in a strictly quantitative manner, of drug effects in man on the basis information from *in vitro* bioassays and/or *in vivo* animal studies [1]. This impels developing tools with much improved properties for extrapolation and prediction, such as mechanism-based PK-PD modelling and simulation

Methods: Mechanism-based PK-PD models are based on principles from systems biology and contain specific expressions to characterize processes on the causal path between plasma concentration and response. This includes a) the target distribution, b) the target interaction/activation and c) transduction and the homeostatic control mechanisms, which may be operative. The utilisation of these models relies on novel biomarkers characterising specific processes on the causal path in a quantitative manner. An essential feature of mechanism-based PK-PD models is the strict distinction between “drug-specific” and “biological system-specific” pharmacodynamic parameters to describe *in vivo* drug effects [2,3,4,5]

The latest development in mechanism-based PK-PD modeling has been the introduction of the concept of disease systems analysis, to characterize drug effects in disease processes and disease progression. Disease systems analysis aims at the distinction between drug effects on the “disease status” *versus* the “disease process” enabling the prediction of long-term treatment effect [6].

Results: We have successfully developed mechanism-based PK-PD models for drugs acting at various targets including A₁ Adenosine, μ Opioid, 5-HT_{1A} Serotonin and GABA_A receptors. Our findings show that in general a drug's *in vivo* intrinsic efficacy can be accurately predicted on the basis of *in vitro* bioassays. Prediction of the *in vivo* potency on the other hand appears to be more difficult, presumably as result of complexities at the level of the target site distribution. Our results also show that equilibrium concentration-effect relationships can be readily scaled from pre-clinical animal models to humans. The utility of this approach has recently been demonstrated for (semi-)synthetic opioids where a mechanism-based PK-PD model has been developed which can predict the clinical analgesic and respiratory depressant effects on the basis of preclinical *in vitro* and *in vivo* data [7]. In contrast, the scaling of transduction and homeostatic feedback mechanisms appears to be more complex. An example of the latter is our work on the allometric scaling of different biomarkers for 5-HT_{1A} receptor agonists from preclinical *in vitro* and *in vivo* models to man [8]. The first application of disease systems analysis has been in the field of type 2 diabetes mellitus, where it has been shown that drug effects on the deterioration of beta cell function and insulin sensitivity can be quantified by analyzing a cascade of biomarker responses [9].

Conclusions: It is concluded that mechanism-based PK-PD models provide a scientific basis for the prediction of efficacy and safety of novel drugs in humans on the basis information from *in vitro* bioassays and/or *in vivo* animal studies.

References:

- [1] Kola I and Landis J (2004) Can the pharmaceutical industry reduce attrition rates? *Nature Rev Drug Discov.* 3, 711-716.
- [2] Danhof M, Van der Graaf PH, Jonker DM, Visser SAG and Zuideveld KP, (2007) Mechanism-based pharmacokinetic-pharmacodynamic modeling for the prediction of *in vivo* drug concentration-effect relationships-Application in drug candidate selection and lead optimization. In: *Comprehensive Medicinal Chemistry II, Part 5*, Testa, B. and Van de Waterbeemd, H. eds., pp. 885-908. Elsevier

- [3] Danhof M, DeJongh J, DeLange ECM, Della Pasqua OE, Ploeger BA and Voskuyl RA (2007) Mechanism-based pharmacokinetic-pharmacodynamic modeling: biophase distribution, receptor theory and dynamical systems analysis. *Ann. Rev. Pharmacol. Toxicol.* 47, 357-400.
- [4] Danhof M, DeLange ECM, Della Pasqua OE, Ploeger BA and Voskuyl RA (2008) Mechanism-based pharmacokinetic-pharmacodynamic (PK-PD) modeling in translational drugs research *Trends in Pharmacol. Sci.* in press.
- [5] Danhof, M, Alvan G., Dahl SG, Kuhlmann J, and Paintaud G. (2005) Mechanism-based pharmacokinetic-pharmacodynamic modeling-a new classification of biomarkers. *Pharm. Res.* 22: 1432-1437.
- [6] Post, TM Freijer JI, DeJongh J and Danhof M (2005) Disease system analysis: basic disease progression models in degenerative disease. *Pharm. Res.* 22: 1038-1049.
- [7] Yassen A, Olofsen E, Kan J, Dahan A and Danhof M (2007) Animal-to-human extrapolation of the pharmacokinetic and pharmacodynamic properties of buprenorphine. *Clin Pharmacokinet.* 46:433-447.
- [8] Zuideveld KP, Van der Graaf PH, Peletier LA and Danhof M (2007) Allometric scaling of pharmacodynamic responses: application to 5-Ht1A receptor mediated responses from rat to man. *Pharm Res.* 24:2031-2039.
- [9] DeWinter W, DeJongh J, Post T, Ploeger B, Urquhart R, Moules I, Eckland D and Danhof M. (2006) A mechanism-based disease progression model for comparison of long-term effects of pioglitazone, metformin and gliclazide on disease processes underlying Type 2 Diabetes Mellitus. *J Pharmacokinet Pharmacodyn.* 33:313-343.